



Silvi, M., & Aggarwal, V. K. (2019). Radical Addition to Strained σ -Bonds Enables the Stereocontrolled Synthesis of Cyclobutyl Boronic Esters. *Journal of the American Chemical Society*, 141(24), 9511-9515. <https://doi.org/10.1021/jacs.9b03653>

Peer reviewed version

Link to published version (if available):
[10.1021/jacs.9b03653](https://doi.org/10.1021/jacs.9b03653)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via ACS at <https://pubs.acs.org/doi/10.1021/jacs.9b03653>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Radical Addition to Strained σ -Bonds Enables the Stereocontrolled Synthesis of Cyclobutyl Boronic Esters

Mattia Silvi, and Varinder K. Aggarwal*

School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, United Kingdom.

Supporting Information Placeholder


ABSTRACT: Whilst radical additions to π -bonds are well established, additions to σ -bonds are far less explored. We have found that electron deficient radicals derived from alkyl iodides under visible light irradiation add to the central strained bond of bicyclobutyl (BCB)-boronate complexes and lead to 1,3-alkyl disubstituted cyclobutyl boronic esters in high yields, with full stereospecificity and high levels of stereoselectivity. Novel cyclobutyl-substituted structures, including peptide and steroid boronic ester derivatives can be accessed. Additionally, although the use of electron-rich alkyl iodides as radical precursors was found to be ineffective, an alternative route involving alkylsulfonylation of the BCB-boronate, followed by reductive desulfonylation provided access to simple alkyl substituted cyclobutane products.

The field of radical chemistry¹ has witnessed a renaissance over the last decade,² triggered largely by the introduction of photoredox methods (electron transfer photocatalysis)³⁻⁴ for the generation of open-shell species under mild conditions. The radicals generated have been utilized in a broad array of reactions including addition to π -systems (e.g. olefins, alkynes, aromatics),²⁻⁴ radical-metal recombination⁵ and radical-radical cross-coupling.⁶

We and others recently showed that electrophilic radicals readily add to π -bonds of unsaturated boronate complexes **1** (Scheme 1a).⁷ Subsequent one-electron oxidation of the resulting intermediate radical anion **2** triggered a 1,2-metallate rearrangement leading to densely functionalized boronic esters **3**. We questioned whether less activated σ -bonds, embedded in a strained molecule,⁸ could undergo analogous radical reactivity as this would significantly enhance the scope of both boron and radical chemistry. In support of this hypothesis, the addition of radicals to exceptionally strained σ -bonds has been observed,⁹ although studies in this area remain limited. Specifically, we sought to add radicals to the central σ -bond of a strained bicyclobutyl (BCB)-boronate complex **4** (Scheme 1b), which we recently showed was capable of reacting with electrophilic palladium(II) intermediates.¹⁰ Following radical addition, one-electron oxidation⁷ would trigger a 1,2-metallate rearrangement¹¹ leading to cyclobutyl boronic esters **6**. Not only does this proposal present novel reactivity between boronate complexes and radicals, but it also offers a mechanistically distinct method to prepare challenging polysubstituted cyclobutanes.¹² Such entities have the potential to open up considerable chemical space due to the three readily diversifiable positions (radical precursor,⁴ boronic ester substituent, and the boron atom¹³ itself). Furthermore,

cyclobutanes are of increasing interest in medicinal chemistry,¹⁴ adding significant motivation for developing this methodology. Herein, we show that radicals can indeed add to the central σ -bond of a strained BCB-boronate complexes, thereby providing a stereocontrolled synthesis of 1,3-dialkyl substituted cyclobutyl boronic esters **6**.


Scheme 1. a) Radical addition to π -bonds of unsaturated boronate complexes. b) Proposed radical addition to σ -bonds of BCB-boronate complexes.



We began our studies by focusing on developing a radical trifluoromethylation process,^{2c} as the trifluoromethyl group is a highly sought-after moiety in medicinal chemistry.¹⁵ For the optimization studies, the BCB-boronate complex **3a** was generated *in situ* by adding *t*-BuLi to a mixture of sulfoxide **7** and cyclohexyl boronic acid pinacol ester **8a** in THF. An excess (5 equiv.) of trifluoromethyl iodide-DMSO complex and 1 mol% of the photocatalyst Ru(bpy)₃²⁺ were then added under blue LED irradiation. After 16 h we were delighted to find that the expected boronic ester **6a** was formed in good yield and moderate diastereoselectivity (Table 1, entry 1). A brief solvent screen showed that THF was superior to DMSO and MeCN (entries 2-3). Performing the reaction at -78 °C¹⁶ was found to significantly improve the process, providing the desired compound **6a** in high yield and as a single diastereoisomer (entry 4; stereochemistry established by ¹H-NMR NOE analysis,

see SI for details). In line with our previous studies with vinyl boronates,^{7b} we found that the photocatalyst was unnecessary for the process (entry 5). Finally, the loading of radical precursor could be decreased to 3 equivalents, and the reaction time shortened to 1 hour, without any loss in reaction efficiency (entry 6). A control experiment highlighted the importance of light, as only traces of **6a** were observed in the absence of blue LED irradiation (entry 7). Finally, performing the reaction in the presence of a radical inhibitor (TEMPO) led to complete inhibition (entry 8), and to the detection of the TEMPO–CF₃ adduct (see SI), which is indicative of the involvement of radical species in the process.

Table 1. Optimization Studies.




Entry ^a	Solvent	T (°C)	6a (%) ^b	d.r. ^c
1 ^d	THF	25	66	5:1
2 ^e	DMSO	25	52	6:1
3 ^e	CH ₃ CN	25	46	5:1
4 ^d	THF	-78	73	>20:1
5	THF	-78	75	>20:1
6 ^f	THF	-78	79 (74)	>20:1
7 ^g	THF	-78 to 25	6	5:1
8 ^{h,f}	THF	-78	0	-

^a All the reactions performed on a 0.1 mmol scale, light source: Kessil blue LED lamp (see SI), reaction time: 16 h. Cy: cyclohexyl.
^b ¹⁹F-NMR yield using trifluorotoluene as internal standard, number in parentheses is the isolated yield of a 0.2 mmol scale reaction.
^c Determined by ¹⁹F-NMR. ^d 1 mol% of Ru(bpy)₃(PF₆)₂ used. ^e 1 mol% of Ru(bpy)₃Cl₂ used. ^f Reaction carried out with 3 equiv. of CF₃I·2DMSO, reaction time: 1 hour. ^g Reaction carried out without blue LED irradiation. ^h Reaction carried out in the presence of 1 equiv. of 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO).

With the optimized conditions in hand, we explored the scope of the radical trifluoromethylation reaction by varying the structure of the starting boronic ester. The reactions were found to occur with high diastereoselectivity, and simple chromatographic separation of the isomers allowed isolation of the products as single diastereoisomers. A variety of alkyl groups could be accommodated on the boronic ester starting material, from a simple Me group up to a *t*-Bu group, showing that the reaction tolerates the full spectrum of steric demand and affords products **6b–d** with high stereoselectivity (Scheme 2). Aromatic boronic esters were also successfully employed, although only moderate levels of diastereoselectivity were achieved (**6e–g**).¹⁷ A cyclobutyl boronic ester was also explored, giving the strained boronic ester **6h**, featuring two adjacent cyclobutyl rings with high diastereoselectivity. Enantioenriched boronic esters featuring a chiral center in the α -position gave **6i–k** with complete retention of stereochemistry. The pyrrolidine boronic ester substrate is especially noteworthy as this is normally a poor migrating group¹⁸ but worked well under our standard conditions giving **6k** in good yield and with high stereocontrol. In addition, cyclobutyl boronic esters bearing complex chiral structures and biologically relevant substituents could be obtained through this methodology with high stereocontrol (**6l–o**).


Scheme 2. Scope of boronic esters for the radical trifluoromethylation reaction^a



^a Unless noted otherwise, isolated yields of single diastereoisomer in reactions carried out under a nitrogen atmosphere, with 1.2 equiv. of **7**, 1.0 equiv. **8** and 3 equiv. of CF₃I·2DMSO, using Schlenk techniques on a 0.2 mmol scale. Diastereomeric ratio measured by ¹⁹F-NMR analysis of the crude reaction mixture and refers to stereochemistry across the cyclobutane ring: the migration was in all cases observed to be stereospecific with respect to the migrating carbon. ^b Reaction time: 24 hours, propionitrile used as solvent. ^c Compound isolated as a mixture of diastereoisomers.

We then sought to explore the scope of radical precursors that could be employed. Longer perfluoroalkyl chains could be used, resulting in similarly high yield and stereoselectivity (Scheme 3, compound **6p**). A number of alkyl halides carrying electron withdrawing groups – including nitriles, esters, primary amides and sulfones – reacted with high yield and moderate to good diastereoselectivity (**6q–t**). The stereoselectivity of the process was observed to follow the trend CN < COOEt < SO₂Ph, indicating that steric factors influence the outcome of the reaction. Remarkably, dipeptide **6u** and tigogenin derivative **6v** could also be efficiently synthesized from radical precursors derived from the corresponding biologically relevant molecule, showcasing the applicability of this process for late stage functionalizations.


Scheme 3. Scope of radical precursors^a



^a Isolated yields of mixtures of diastereoisomers in reactions carried out under a nitrogen atmosphere, with 1.2 equiv. of **7**, 1.0 equiv. **8a** and 2 equiv. of iodide **5**, using Schlenk techniques on a 0.2 mmol scale. Diastereomeric ratio measured by ¹H-NMR analysis of the crude reaction mixture. ^b Ru(bpy)₃(PF₆)₂ (1 mol%) was used. ^c Reaction time: 24 hours.

Unfortunately, non-activated alkyl iodides (e.g. benzyl iodide, methyl iodide) were found to be unreactive in the process, presumably because the polarity of the electron rich BCB-boronate, limiting the scope of the methodology. To overcome this limitation, we considered the use of sulfones as traceless activating groups.¹⁹ Pleasingly, sequential radical alkylsulfonylation followed by reductive cleavage of the sulfone enabled access to alkylated products **9a** and **9b** in high yields (Scheme 4). Interestingly, the cyclobutylmethyl radical intermediate involved in the desulfonylation did not undergo ring opening [*k*=4.5·10² s⁻¹],²⁰ showing that reduction/protonation are faster processes.


Scheme 4. Strategy to introduce simple alkyl groups



The synthetic utility of the products was demonstrated by carrying out a number of functional group interconversions of the


boronic ester. For example, cyclobutane **6a** underwent stereospecific oxidation to alcohol **10a**, conversion to potassium trifluoroborate salt²¹ **10b**, Zweifel olefination²² to give alkene **10c**, and Matteson homologation²³ to afford boronic ester **10d** (Scheme 5).

Scheme 5. Product functionalization reactions



The mechanism of this photochemical radical reaction of BCB-boronate complexes likely bears similarities to that previously reported for vinyl boronate complexes.^{7b} We propose that the light sensitive alkyl iodide undergoes photolytic initiation generating radical **11**, which adds to the strained central σ-bond of BCB-boronate **3** leading to the electron-rich radical anion **12** (Scheme 6). Single electron transfer to another molecule of alkyl iodide **5** regenerates radical **11** and forms zwitterionic species **13**, which undergoes fast 1,2-metallate rearrangement to give the final product **6**.²⁴ We believe that the single electron-transfer step is also the stereoselectivity determining step. As depicted in Scheme 6, alkyl iodide **5** approaches radical intermediate **12** from the less hindered face of the cyclobutane ring, away from the alkyl iodide-derived substituent. As the electron is transferred, 1,2-migration occurs on the opposite lobe of the p orbital before the molecule has time to undergo bond rotation around the C⁺–B bond. This accounts for the *cis*-relationship between the two alkyl groups in **6**.

Scheme 6. Proposed reaction mechanism



In conclusion, we have shown that bicyclobutyl boronate complexes are excellent radical traps for electrophilic radicals, providing a novel visible light-mediated three-component coupling. Furthermore, the reaction can be conducted in the absence of photocatalysts. The reactions occur with moderate to excellent diastereoselectivity, are fully stereospecific, and allow the synthesis of complex chiral cyclobutanes, including peptide and steroid derivatives. The reactivity is induced by strain release-driven radical

addition to σ -bonds, a pathway that will no doubt find further application in the synthesis of cyclobutyl-containing natural products and drug candidates

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

*v.aggarwal@bristol.ac.uk

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

M.S. thanks the EU for H-2020 Marie Skłodowska-Curie Fellowship (grant no. 744242) and EPSRC (EP/R012695/1) for financial support. We thank Dr. Adam Noble for proofreading this manuscript.

REFERENCES

- (1) a) *Encyclopedia of Radicals in Chemistry, Biology and Materials*; Studer, A., Chatgililoglu, C., Eds.; Wiley: 2012. b) *Radicals in Organic Synthesis*; Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, Germany, 2001.
- (2) For selected reviews: a) Studer, A.; Curran, D. P. Organocatalysis and C-H Activation Meet Radical- and Electron-Transfer Reactions. *Angew. Chem. Int. Ed.* **2011**, *50*, 5018. b) Finkbeiner, P.; Nachtsheim, B. J. Iodine in Modern Oxidation Catalysis. *Synthesis* **2013**, *45*, 979. c) Studer, A. A "Renaissance" in Radical Trifluoromethylation. *Angew. Chem. Int. Ed.* **2012**, *51*, 8950. d) Jahn, U. Radicals in Transition Metal Catalyzed Reactions? Transition Metal Catalyzed Radical Reactions?: A Fruitful Interplay Anyway. In *Radicals in Synthesis III* Heinrich, M., Gansäuer, A., Eds.; Springer: Berlin, Heidelberg, 2012; pp 323–451. e) Sibi, M. P.; Landais, Y. Csp³-F Bond Formation: A Free-Radical Approach. *Angew. Chem. Int. Ed.* **2013**, *52*, 3570. f) Romero, K. J.; Galliher, M. S.; Pratt, D. A.; Stephenson, C. R. J. Radicals in Natural Product Synthesis. *Chem. Soc. Rev.* **2018**, *47*, 7851. g) Studer, A.; Curran, D. P. Catalysis of Radical Reactions: A Radical Chemistry Perspective. *Angew. Chem. Int. Ed.* **2016**, *55*, 58. h) Xiong, T.; Zhang, Q. New Amination Strategies Based on Nitrogen-Centered Radical Chemistry. *Chem. Soc. Rev.* **2016**, *45*, 3069.
- (3) a) Nicewicz, D. A.; MacMillan, D. W. C. Merging Photoredox Catalysis with Organocatalysis: The Direct Asymmetric Alkylation of Aldehydes. *Science* **2008**, *322*, 77. b) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. Efficient Visible Light Photocatalysis of [2+2] Enone Cycloadditions. *J. Am. Chem. Soc.* **2008**, *130*, 12886. c) Narayanam, J. M. R.; Tucker, J. W.; Stephenson, C. R. J. Electron-Transfer Photoredox Catalysis: Development of a Tin-Free Reductive Dehalogenation Reaction. *J. Am. Chem. Soc.* **2009**, *131*, 8756.
- (4) a) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322. b) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81*, 6898. c) Goddard, J.-P.; Ollivier, C.; Fensterbank, L. Photoredox Catalysis for the Generation of Carbon Centered Radicals. *Acc. Chem. Res.* **2016**, *49*, 1924. d) Staveness, D.; Bosque, I.; Stephenson, C. R. J. Free Radical Chemistry Enabled by Visible Light-Induced Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 2295.
- (5) a) Kalyani, D.; McMurtrey, K. B.; Neufeldt, S. R.; Sanford, M. S. Room-Temperature C–H Arylation: Merger of Pd-Catalyzed C–H Functionalization and Visible-Light Photocatalysis. *J. Am. Chem. Soc.* **2011**, *133*, 18566. b) Zuo, Z.; Ahneman, D. T.; Chu, L.; Terrett, J. A.; Doyle, A. G.; MacMillan, D. W. C. Merging Photoredox with Nickel Catalysis: Coupling of α -Carboxyl sp³-Carbons with Aryl Halides. *Science* **2014**, *345*, 437. c) Sahoo, B.; Hopkinson, M. N.; Glorius, F. Combining Gold and Photoredox Catalysis: Visible Light-Mediated Oxy- and Aminoarylation of Alkenes. *J. Am. Chem. Soc.* **2013**, *135*, 5505. d) Ye, Y.; Sanford, M. S. Merging Visible-Light Photocatalysis and Transition-Metal Catalysis in the Copper-Catalyzed Trifluoromethylation of Boronic Acids with CF₃I. *J. Am. Chem. Soc.* **2012**, *134*, 9034. e) Twilton, J.; Le, C.; Zhang, P.; Shaw, M. H.; Evans, R. W.; MacMillan, D. W. C. The merger of transition metal and photocatalysis. *Nat. Rev. Chem.* **2017**, *1*, 0052.
- (6) For selected examples: a) McNally, A.; Prier, C. K.; MacMillan, D. W. C. Discovery of an α -Amino C–H Arylation Reaction Using the Strategy of Accelerated Serendipity. *Science* **2011**, *334*, 1114. b) Cuthbertson, J. D.; MacMillan, D. W. C. The Direct Arylation of Allylic sp³ C–H Bonds via Organic and Photoredox Catalysis. *Nature* **2015**, *519*, 74. c) Pirmot, M. T.; Rankic, D. A.; Martin, D. B. C.; MacMillan, D. W. C. Photoredox Activation for the Direct β -Arylation of Ketones and Aldehydes. *Science* **2013**, *339*, 1593.
- (7) a) Kischkewitz, M.; Okamoto, K.; Mück-Lichtenfeld, C.; Studer, A. Radical-polar crossover reactions of vinylboron ate complexes. *Science* **2017**, *355*, 936. b) Silvi, M.; Sandford, C.; Aggarwal, V. K. Merging Photoredox with 1,2-Metallate Rearrangements: The Photochemical Alkylation of Vinyl Boronate Complexes. *J. Am. Chem. Soc.* **2017**, *139*, 5736. c) Lovinger, G. J.; Morken, J. P. Ni-Catalyzed Enantioselective Conjugative Coupling with C(sp³) Electrophiles: A Radical-Ionic Mechanistic Dichotomy. *J. Am. Chem. Soc.* **2017**, *139*, 17293. d) Gerleve, C.; Kischkewitz, M.; Studer, A. Synthesis of α -Chiral Ketones and Chiral Alkanes Using Radical Polar Crossover Reactions of Vinyl Boron Ate Complexes. *Angew. Chem. Int. Ed.* **2018**, *57*, 2441. e) Silvi, M.; Schrof, R.; Noble, A.; Aggarwal, V. K. Enantiospecific Three-Component Alkylation of Furan and Indole. *Chem. Eur. J.* **2018**, *24*, 4279. f) Tappin, N. D. C.; Gnägi-Lux, M.; Renaud, P. Radical-Triggered Three-Component Coupling Reaction of Alkenylboronates, α -Halocarbonyl Compounds, and Organolithium Reagents: The Inverse Ylid Mechanism. *Chem. Eur. J.* **2018**, *24*, 11498.
- (8) Recent reports of utilization of strained systems: a) Gianatassio, R.; Lopchuk, J. M.; Wang, J.; Pan, C.-M.; Malins, L. R.; Prieto, L.; Brandt, T. A.; Collins, M. R.; Gallego, G. M.; Sach, N. W.; et al. Strain-release amination. *Science* **2016**, *351*, 241. b) Lopchuk, J. M.; Fjelbye, K.; Kawamata, Y.; Malins, L. R.; Pan, C.-M.; Gianatassio, R.; Wang, J.; Prieto, L.; Bradow, J.; Brandt, T. A.; et al. Strain-Release Heteroatom Functionalization: Development, Scope, and Stereospecificity. *J. Am. Chem. Soc.* **2017**, *139*, 3209. c) Makarov, I. S.; Brocklehurst, C. E.; Karaghiosoff, K.; Koch, G.; Knochel, P. Synthesis of Bicyclo[1.1.1]pentane Bioisosteres of Internal Alkynes and para-Disubstituted Benzenes from [1.1.1]Propellane. *Angew. Chem. Int. Ed.* **2017**, *56*, 12774.
- (9) a) Wiberg, K. B. Small ring propellanes. *Chem. Rev.* **1989**, *89*, 975. b) Della, E. W.; Lochert, I. J. Synthesis of Bridgehead-Substituted Bicyclo[1.1.1]Pentanes. A Review. *Org. Prep. Proc. Int.* **1996**, *28*, 411. c) Levin, M. D.; Kaszynski, P.; Michl, J. Bicyclo[1.1.1]pentanes, [n]Staffanes, [1.1.1]Propellanes, and Tricyclo[2.1.0.0^{2,5}]pentanes. *Chem. Rev.* **2000**, *100*, 169. d) Wiberg, K. B.; Waddell, S. T.; Laidig, K. [1.1.1]Propellane: Reaction with free radicals. *Tetrahedron Lett.* **1986**, *27*, 1553. e) Bunker, K. D.; Sach, N. W.; Huang, Q.; Richardson, P. F. Scalable Synthesis of 1-Bicyclo[1.1.1]pentylamine via a Hydrohydrazination Reaction. *Org. Lett.* **2011**, *13*, 4746. f) Kanazawa, J.; Maeda, K.; Uchiyama, M. Radical Multicomponent Carboamination of [1.1.1]Propellane. *J. Am. Chem. Soc.* **2017**, *139*, 17791. g) Caputo, D. F. J.; Arroniz, C.; Dürr, A. B.; Mousseau, J. J.; Stepan, A. F.; Mansfield, S. J.; Anderson, E. A. Synthesis and Applications of Highly Functionalized 1-Halo-3-Substituted Bicyclo[1.1.1]pentanes. *Chem. Sci.* **2018**, *9*, 5295. h) Hoz, S. Bicyclo[1.1.0]butane. In *The Chemistry of the Cyclopropyl Group. Volume 1*; Rappoport, Z., Ed.; 1987; pp 1158–1160. i) Wu, X.; Hao, W.; Ye, K.-Y.; Jiang, B.; Pombar, G.; Song, Z.; Lin, S. Ti-Catalyzed Radical Alkylation of Secondary and Tertiary Alkyl Chlorides Using Michael Acceptors. *J. Am. Chem. Soc.* **2018**, *140*, 14836.
- (10) Fawcett, A.; Biberger, T.; Aggarwal, V. K. Carbopalladation of C–C σ -Bonds Enabled by Strained Boronate Complexes. *Nat. Chem.* **2019**, *11*, 117.
- (11) a) Matteson, D. S.; Mah, R. W. H. Neighboring Boron in Nucleophilic Displacement. *J. Am. Chem. Soc.* **1963**, *85*, 2599. b) Matteson, D. S. Boronic Esters in Asymmetric Synthesis. *J. Org. Chem.* **2013**, *78*, 10009. c) Leonori, D.; Aggarwal, V. K. Lithiation–Borylation Methodology and Its Application in Synthesis. *Acc. Chem. Res.* **2014**, *47*, 3174. d) Thomas, S. P.; French, R. M.; Jheengut, V.; Aggarwal, V. K. Homologation and Alkylation of Boronic Esters and Boranes by 1,2-Metallate Rearrangement of Boron Ate Complexes. *Chem. Rev.* **2009**, *9*, 24.
- (12) a) Lee-Ruff, E. Synthesis of Cyclobutanes. In *The Chemistry of Cyclobutanes*; Rappoport, Z.; Liebman, J. F., Eds.; 2005; pp 281–355. b)

Poplata, S.; Tröster, A.; Zou, Y.-Q.; Bach, T. Recent Advances in the Synthesis of Cyclobutanes by Olefin [2+2] Photocycloaddition Reactions. *Chem. Rev.* **2016**, *116*, 9748. c) Hoffmann, N. Photochemical Reactions as Key Steps in Organic Synthesis. *Chem. Rev.* **2008**, *108*, 1052. d) Lee-Ruff, E.; Mladenova, G. Enantiomerically Pure Cyclobutane Derivatives and Their Use in Organic Synthesis. *Chem. Rev.* **2003**, *103*, 1449.

(13) Sandford, C.; Aggarwal, V. K. Stereospecific Functionalizations and Transformations of Secondary and Tertiary Boronic Esters. *Chem. Commun.* **2017**, *53*, 5481.

(14) a) Dembitsky, V. M. Naturally Occurring Bioactive Cyclobutane-Containing (CBC) Alkaloids in Fungi, Fungal Endophytes, and Plants. *Phytomedicine* **2014**, *21*, 1559. b) Blakemore, D. C.; Bryans, J. S.; Carnell, P.; Carr, C. L.; Chessum, N. E. A.; Field, M. J.; Kinsella, N.; Osborne, S. A.; Warren, A. N.; Williams, S. C. Synthesis and In Vivo Evaluation of Bicyclic Gababutins. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 461. c) Slade, J.; Bajwa, J.; Liu, H.; Parker, D.; Vivel, J.; Chen, G.-P.; Calieni, J.; Villhauer, E.; Prasad, K.; Repić, O.; et al. A Concise Synthesis of a Novel Insulin-Like Growth Factor I Receptor (IGF-IR) Inhibitor. *Org. Proc. Res. Dev.* **2007**, *11*, 825. d) Wroblewski, M. L.; Reichard, G. A.; Paliwal, S.; Shah, S.; Tsui, H.-C.; Duffy, R. A.; Lachowicz, J. E.; Morgan, C. A.; Varty, G. B.; Shih, N.-Y. Cyclobutane derivatives as potent NK1 selective antagonists. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3859.

(15) a) Müller, K.; Faeh, C.; Diederich, F. Fluorine in Pharmaceuticals: Looking Beyond Intuition. *Science* **2007**, *317*, 1881. b) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. Fluorine in Medicinal Chemistry. *Chem. Soc. Rev.* **2008**, *37*, 320. c) Hagmann, W. K. The Many Roles for Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2008**, *51*, 4359.

(16) Illumination at low temperature could be effectively, practically and reproducibly achieved by using a commercially available high-power LED light source, a cryostat and a mirror-walled dewar. For experimental details and a picture of the reaction set-up, see Supporting Information.

(17) We believe that the lower stereoselectivity observed for aromatic boronic esters, when compared to alkyl boronic esters, results from these substrates reacting via a different mechanism, in which the radical anion

intermediate can undergo 1,2-aryl migration prior to single electron oxidation. This is discussed in more detail in the Supporting Information.

(18) Varela, A.; Garve, L. K. B.; Leonori, D.; Aggarwal, V. K. Stereoccontrolled Total Synthesis of (–)-Stemaphylline. *Angew. Chem. Int. Ed.* **2017**, *56*, 2127.

(19) a) Gui, J.; Zhou, Q.; Pan, C.-M.; Yabe, Y.; Burns, A. C.; Collins, M. R.; Ornelas, M. A.; Ishihara, Y.; Baran, P. S. C–H Methylation of Heteroarenes Inspired by Radical SAM Methyl Transferase. *J. Am. Chem. Soc.* **2014**, *136*, 4853. b) Filippini, G.; Silvi, M.; Melchiorre, P. Enantioselective Formal α -Methylation and α -Benzoylation of Aldehydes by Means of Photo-organocatalysis. *Angew. Chem. Int. Ed.* **2017**, *56*, 4447.

(20) Beckwith, A. L. J.; Moad, G. The Kinetics and Mechanism of Ring Opening of Radicals Containing the Cyclobutylcarbinyl System. *J. Chem. Soc., Perkin Trans. 2* **1980**, 1083.

(21) Bagutski, V.; Ros, A.; Aggarwal, V. K. Improved Method for the Conversion of Pinacolboronic Esters into Trifluoroborate Salts: Facile Synthesis of Chiral Secondary and Tertiary Trifluoroborates. *Tetrahedron* **2009**, *65*, 9956.

(22) a) Zweifel, G.; Arzoumanian, H.; Whitney, C. C. A Convenient Stereoselective Synthesis of Substituted Alkenes via Hydroboration-Iodination of Alkynes. *J. Am. Chem. Soc.* **1967**, *89*, 3652. b) Armstrong, R. J.; Aggarwal, V. K. 50 Years of Zweifel Olefination: A Transition-Metal-Free Coupling *Synthesis* **2017**, *49*, 3323.

(23) a) Sadhu, K. M.; Matteson, D. S. Chloromethylolithium: Efficient Generation and Capture by Boronic Esters and a Simple Preparation of Diisopropyl (Chloromethyl)Boronate. *Organometallics* **1985**, *4*, 1687. b) Balieu, S.; Hallett, G. E.; Burns, M.; Bootwicha, T.; Studley, J.; Aggarwal, V. K. Toward Ideality: The Synthesis of (+)-Kalkitoxin and (+)-Hydroxyphthioceranic Acid by Assembly-Line Synthesis *J. Am. Chem. Soc.* **2015**, *137*, 4398.

(24) Presently, we cannot exclude the possibility of an atom transfer mechanism. In this scenario, radical anion **12** abstracts an I atom from another molecule of **5**, generating radical **11** and an α -iodo boronate complex, which is expected to undergo 1,2-migration rapidly to afford product **6** (see ref. 7).

Insert Table of Contents artwork here

